



Factors affecting denitrification rates in soils




In grazed pastures urine patches are the main sources of nitrous oxide emissions and nitrate leaching



Nitrate leaching is greatest below the animal urine patch



N_2O emissions from urine amended soil



Source: Sherlock et al. (unpubl. data)

How can we mitigate N₂O emissions from excreta N?

- Change the animals diet?
 - modify the C/N ratio?
- Stand-off/feed pads?

Covered feed & loafing pad



(Cecile De Klein, AgResearch 2008)

Uncovered feed & loafing pad




(Cecile De Klein, AgResearch 2008)




How can we mitigate excreta N?

- Change the animals diet?
 - Increase C and modify the C/N ratio?
 - Stand-off/feed pads
 - Use nitrification inhibitors e.g. dicyandiamide (DCD)


DCD reduces the activity of the *Nitrosomonas* bacteria in the soil (Bacteriostatic effect)




The inhibitor temporarily blocks the active site of a specific enzyme (ammonia monoxygenase)




Recommend DCD is applied in May and August because most losses occur between late autumn and early spring




DCD reduced N₂O emissions by 73% on Templeton soil in Canterbury, NZ (Di et al., 2007).




N₂O emissions reduced by 67% in the Canterbury Lismore soil (Di et al., 2007)






N₂O emissions reduced by 61% in the Waikato Horotiu soil (Di et al., 2007)



N₂O emissions reduced by 69% in the Taupo pumice soil lysimeters (Di et al., 2007)





Internationally peer reviewed and published literature:

- Di, H.J., Cameron, K.C., 2002. The use of a nitrification inhibitor, dicyandiamide (DCD), to decrease nitrate leaching and nitrous oxide emissions in a simulated grazed and irrigated grassland. *Soil Use and Management* 18, 395-403.
- Di, H.J., Cameron, K.C., 2003. Mitigation of nitrous oxide emissions in spray - irrigated grassland by treating the soil with dicyandiamide, a nitrification inhibitor. *Soil Use and Management* 19, 284-290.
- Di, H.J., Cameron, K.C., 2004a. Effects of the nitrification inhibitor dicyandiamide on potassium, magnesium and calcium leaching in grazed grassland. *Soil Use & Management* 20, 2-7.
- Di, H.J., Cameron, K.C., 2004b. Treating pasture soil with a nitrification inhibitor, eco-nTM, to decrease nitrate leaching in a deep sandy soil under spray irrigation - a lysimeter study. *New Zealand Journal of Agricultural Research* 47, 351-361.
- Di, H.J., Cameron, K.C., 2004c. Effects of temperature and application rate of a nitrification inhibitor, dicyandiamide (DCD), on nitrification rate and microbial biomass in a grazed pasture soil. *Australian Journal of Soil Research* 42, 927-932.
- Di, H.J., Cameron, K.C., 2005. Reducing environmental impacts of agriculture by using a fine particle suspension nitrification inhibitor to decrease nitrate leaching from grazed pastures. *Agriculture, Ecosystems and Environment* 109, 202-212.
- Di, H.J., Cameron, K.C., 2006. Nitrous oxide emissions from two dairy pasture soils as affected by different rates of a fine particle suspension nitrification inhibitor, dicyandiamide. *Biology and Fertility of Soils* 42, 472-480.

Some of the data required to develop an N₂O inventory consists of:

- Animal numbers
- N excreta
- Direct emission factors
 - EF1 N₂O emitted from fertiliser
 - EF3_{PRP} N₂O emitted from urine and dung
- Indirect emission factors
 - EF5 N₂O emitted from leached N

2006 IPCC guidelines for National Greenhouse Gas Inventories

For a mitigation to be included in an inventory we also must know:

- Animal numbers affected by mitigation tool?
- Changes in N excreted?
- Time period of the year mitigation tool works?
- Changes in direct emission factors ?
- Changes in indirect emission factors ?
- Verification?

Summary of international peer reviewed literature: DCD effect on direct N₂O emissions

- The **reduction** in direct N₂O emissions EF_{3(PRP)} **averaged 67% ± 6%** (std dev. n=4.)

Summary of international peer reviewed literature: DCD effect on NO₃⁻ leaching

- The **reduction** in nitrate leaching
Frac_(LEACH) **averaged 74% ± 4%** (std dev. n=5.)

Calculating DCD emission factors (EF)

$$EF1' \text{ plus DCD} = (IPCC \text{ default } EF1) - (IPCC \text{ default } EF1 \times \frac{50\%}{100\%} \times \frac{5 \text{ months}}{12 \text{ months}}) = 0.0099 \text{ kg N}_2\text{O/kg fertiliser-N}$$

$$EF3_{PRP}' \text{ plus DCD} = (IPCC \text{ default } EF3_{PRP}) - (IPCC \text{ default } EF3_{PRP} \times \frac{50\%}{100\%} \times \frac{5 \text{ months}}{12 \text{ months}}) = 0.0079 \text{ kg N}_2\text{O/kg excreta-N}$$


(Clough et al. 2007 Nutr. Cycl. Agroecosyst. 78:1-14.)

Table 8

Module Submodule Worksheet Sheet	2003 Agriculture (New Zealand) Agricultural soils 4.5 (3 of 5) Direct nitrous oxide emissions from animal production (grazing animals)			
Pasture, range and paddock AWMS	N excretion for AWMS PRP (kg N)	Emission factor for AWMS (EF ₃ PRP) (kg N ₂ O-N/kg N)	Total direct animal prodn. emissions of N ₂ O-N (Gg)	Total direct animal prodn. emissions of N ₂ O (Gg)
PRP nil DCD	1,386,897,313	0.0100	13,869	21,794
PRP plus DCD	143,375,290	0.0079	0,836	1,314

#Assumes 25% of dairy cattle under DCD regime

Effect of DCD on N₂O inventory



Verification ???



DCD application map – providing proof of placement


Application Map Report Field Name: RANG DN 1 Date: 7/7/2004 10:18

Customer details

Date and time of application



Paddock details


Customer Name:	Product Name: eco-n
Field Name: RANG DN 1	Rate: RateData
Note:	Swath: 24.00 M 78.7 feet
	Area Covered: 5.02 ha 12.4 acres



Summary

- Identify main N₂O sources and mitigation strategy.
- Peer reviewed science of mitigation tool.
 - Changes in N excreted?
 - Duration of mitigation effect(s)?
 - Changes in direct emission factors ?
 - Changes in indirect emission factors ?
- Verification.
 - Animal numbers affected by mitigation tool?
 - Land area affected by mitigation tool - Use technology! e.g. GPS.






DCD degradation and microbiology

- Soil bacteria utilise and degrade DCD.
- Numbers of nitrifiers not affected.
- N₂-fixers not affected by DCD.
- Repeat application of DCD in laboratory studies has little or no effect on rate of DCD decomposition or ability of DCD to inhibit nitrification.
- Moir et al. (2007, Soil Use & Mgt) shows no change in response to DCD after four years of field data collection in NZ.

DCD degradation and soil temperature




Kelliher & Clough et al. 2008

Internationally peer reviewed and published literature:

- Di, H.J., Cameron, K.C., 2002. The use of a nitrification inhibitor, dicyandiamide (DCD), to decrease nitrate leaching and nitrous oxide emissions in a simulated grazed and irrigated grassland. *Soil Use and Management* 18, 395-403.
- Di, H.J., Cameron, K.C., 2003. Mitigation of nitrous oxide emissions in spray - irrigated grassland by treating the soil with dicyandiamide, a nitrification inhibitor. *Soil Use and Management* 19, 284-290.
- Di, H.J., Cameron, K.C., 2004a. Effects of the nitrification inhibitor dicyandiamide on potassium, magnesium and calcium leaching in grazed grassland. *Soil Use & Management* 20, 2-7.
- Di, H.J., Cameron, K.C., 2004b. Treating pasture soil with a nitrification inhibitor, eco-nTM, to decrease nitrate leaching in a deep sandy soil under spray irrigation - a lysimeter study. *New Zealand Journal of Agricultural Research* 47, 351-361.
- Di, H.J., Cameron, K.C., 2004c. Effects of temperature and application rate of a nitrification inhibitor, dicyandiamide (DCD), on nitrification rate and microbial biomass in a grazed pasture soil. *Australian Journal of Soil Research* 42, 927-932.
- Di, H.J., Cameron, K.C., 2005. Reducing environmental impacts of agriculture by using a fine particle suspension nitrification inhibitor to decrease nitrate leaching from grazed pastures. *Agriculture, Ecosystems and Environment* 109, 202-212.
- Di, H.J., Cameron, K.C., 2006. Nitrous oxide emissions from two dairy pasture soils as affected by different rates of a fine particle suspension nitrification inhibitor, dicyandiamide. *Biology and Fertility of Soils* 42, 472-480.




Eco-n distribution.



Eco-n supplies over 400 particles per square centimetre.

Scale – 20 cm x 20 cm = 400 cm²

Granular product distribution.



Typically total surface area covered is less than 1% at 100kg per ha.

Scale – 20 cm x 20 cm = 400 cm²

Nitrosomonas bacteria are widespread throughout the soil and they all need to be treated.

